

Formulas as Types

by Khalil Ghorbal (Inria, France)

on November 2020

Cartesian closed categories

» Product


Let \mathcal{A} be a category and $A, B \in \mathcal{A}$. A **product** of A and B consists of an object P and maps π, π' called **projections**

$$A \xleftarrow{\pi} P \xrightarrow{\pi'} B$$

such that for all triples (X, f, g) satisfying

$$A \xleftarrow{f} X \xrightarrow{g} B$$

there is a unique map $p : X \rightarrow P$ making the following diagram commute



» Product

Remarks

- * P is often denoted by $A \times B$
- * p is denoted either by $\langle f, g \rangle$ or $f \times g$
- * **Products do not always exist!**
- * When a product exists, it induces a product-like operation on arrows
- * One can make sense of **the product of zero elements**. It is a terminal object! denoted 1 for convenience.

» Cartesian category

A category \mathcal{A} is **cartesian** if it has finite products (including the product of zero elements 1). That is for every $A, B \in \mathcal{A}$, the product $A \times B$ exists.

» Cartesian closed category

Let \mathcal{A} be a cartesian category. For every object $B \in \mathcal{A}$, we define a functor $- \times B : \mathcal{A} \rightarrow \mathcal{A}$ mapping object A to $A \times B$, and

$$A \xrightarrow{f} A' \quad \mapsto \quad A \times B \xrightarrow{\langle f, 1_B \rangle} A' \times B$$

Cartesian closed category

A category \mathcal{A} is **cartesian closed** if it is cartesian and for each $B \in \mathcal{A}$, the functor $- \times B : \mathcal{A} \rightarrow \mathcal{A}$ has a right adjoint.

We write the right adjoint as $(-)^B$, and, for $C \in \mathcal{A}$, call C^B an **exponential**.

So cartesian closed categories are those **categories with products and exponentials**.

» Important correspondences

In any **cartesian closed** category \mathcal{A} , and $A, B, C \in \mathcal{A}$, by definition of adjunctions

$$\mathcal{A}(A \times B, C) \cong \mathcal{A}(A, C^B)$$

that is arrows $A \times B \rightarrow C$ are in one-to-one correspondence with arrows $A \rightarrow C^B$. We called such operation a **transposition** and denoted it by a bar in both directions. We can also prove ($A = 1$)

$$\mathcal{A}(B, C) \cong \mathcal{A}(1, B^C)$$

so that to each $B \xrightarrow{f} C$ corresponds $1 \xrightarrow{\overline{f\pi'}} B^C$ and to each $1 \xrightarrow{g} B^C$ corresponds $B \xrightarrow{\varepsilon_C \circ \langle g \circ t_B, 1_B \rangle} C$ where $B \xrightarrow{t_B} 1$.

» Higher-order arithmetic

Looks familiar?

In any **cartesian** category \mathcal{A} , and $A, B, C \in \mathcal{A}$

$$A \times 1 \cong A, \quad A \times B \cong B \times A, \quad (A \times B) \times C \cong A \times (B \times C)$$

In any **cartesian closed** category \mathcal{A} , and $A, B, C \in \mathcal{A}$

$$A^1 \cong A, \quad 1^A \cong 1, \quad (A \times B)^C \cong A^C \times B^C, \quad A^{B \times C} \cong (A^C)^B$$

» A category of cartesian closed categories

A **cartesian functor** $F: \mathcal{A} \rightarrow \mathcal{B}$ is a functor that preserves the cartesian closed structure

- * $F(A \times B) = F(A) \times F(B)$
- * $F(\pi_{A,B}) = \pi_{F(A),F(B)}$
- * $F(A^B) = F(A)^{F(B)}$
- * $F(\langle f, g \rangle) = \langle F(f), F(g) \rangle$
- * ...

This defines a category Cart of cartesian closed categories.

» Monad

A **monad** on a category \mathcal{A} is a triple (T, η, μ) where $T: \mathcal{A} \rightarrow \mathcal{A}$

is a functor equipped with a **unit** $\mathcal{A} \begin{matrix} \xrightarrow{1_{\mathcal{A}}} \\ \Downarrow \eta \\ \xrightarrow{T} \end{matrix} \mathcal{A}$

and a **multiplication** $\mathcal{A} \begin{matrix} \xrightarrow{T \circ T} \\ \Downarrow \mu \\ \xrightarrow{T} \end{matrix} \mathcal{A}$

satisfying the associativity and unit laws. That is such that the following diagrams commute:

$$\begin{array}{ccc} T \circ T \circ T & \xrightarrow{T\mu} & T \circ T \\ \mu T \downarrow & & \downarrow \mu \\ T \circ T & \xrightarrow{\mu} & T \end{array}$$

$$\begin{array}{ccc} T & \xrightarrow{T\eta} & T \circ T \\ & \searrow 1_T & \downarrow \mu \\ & & T \end{array}$$

$$\begin{array}{ccc} T & \xrightarrow{\eta T} & T \circ T \\ & \searrow 1_T & \downarrow \mu \\ & & T \end{array}$$


Every adjunction defines a monad!

» Kleisli category

The **Kleisli category** \mathcal{A}_T of a monad (T, η, μ) on \mathcal{A} is a category with

- * the same objects as \mathcal{A}
- * with morphisms $A \rightarrow B$ whenever $A \rightarrow TB$ is a morphism in \mathcal{A}

The identity arrow $1_A : A \rightarrow A$ is defined as $\eta_A : A \rightarrow TA$.
 Two morphisms $f : A \rightarrow B$ and $g : B \rightarrow C$ are composed as $\mu_C \circ Tg \circ f$:



» Natural numbers objects

A **natural numbers object** (or system) in a cartesian closed category \mathcal{A} is an object N and two maps

$$1 \xrightarrow{0} N \xrightarrow{s} N$$

satisfying the following universal property: for any diagram

$$1 \xrightarrow{a} A \xrightarrow{f} A$$

there is a **unique** arrow $N \xrightarrow{h} A$ such that $h \circ 0 = a$ and $h \circ s = f \circ h$. That is such the following diagram commutes

$$\begin{array}{ccccc} 1 & \xrightarrow{0} & N & \xrightarrow{s} & N \\ & \searrow a & \downarrow h & & \downarrow h \\ & & A & \xrightarrow{f} & A \end{array}$$

Drop uniqueness of h to get **weak natural numbers object**.

Deductive systems

» Deductive system

A **deductive system** is a category without the associativity and identity laws axioms.

- * objects are called **formulas**
- * arrows are called **proofs**

» Conjunction calculus

A **conjunction calculus** is a deductive system with

- ⊤ a formula ⊤ (called “true”) such that there is an arrow $A \bullet : A \rightarrow \top$ for each object (a **terminal-like** object ... but we don't have a category)
- ∧ a binary operation ∧ between formulas (called “conjunction”) together with two arrows $A \wedge B \xrightarrow{\pi_{A,B}} A$ and $A \wedge B \xrightarrow{\pi'_{A,B}} B$ inducing a *pairing* of arrows with the same domain often presented as an **inference rule**

$$\frac{C \xrightarrow{f} A \quad C \xrightarrow{g} B}{C \xrightarrow{\langle f, g \rangle} A \wedge B}$$

(a **product-like** construction)

» Proof calculus

proving means *constructing* new proofs (arrows) from a formula (assumption) to another formula (result)

For instance, in conjunction calculus, \wedge is commutative and associative (the labels on arrows are the proofs)

$$* A \wedge B \xrightarrow{\langle \pi'_{A,B}, \pi_{A,B} \rangle} B \wedge A$$

$$* (A \wedge B) \wedge C \xrightarrow{\alpha_{A,B,C}} A \wedge (B \wedge C) \text{ where}$$
$$\alpha_{A,B,C} = \langle \pi_{A,B} \circ \pi_{A \wedge B, C}, \langle \pi'_{A,B} \circ \pi_{A \wedge B, C}, \pi'_{A \wedge B, C} \rangle \rangle$$

» Proof calculus

Inference rules define a **calculus over proofs**: for instance conjunction of formulas (\wedge) induces an operation over arrows (pairing).

Other operations on proofs can be defined out of known ones (derived rules).

For instance

$$\frac{A \xrightarrow{f} B \quad C \xrightarrow{g} D}{A \wedge C \xrightarrow{\langle f \circ \pi_{A,C}, g \circ \pi'_{A,C} \rangle} B \wedge D}$$

defines a “conjunction” on proofs:

$$f \wedge g := \langle f \circ \pi_{A,C}, g \circ \pi'_{A,C} \rangle$$

» Positive intuitionistic propositional calculus

An **positive intuitionistic propositional calculus** is a conjunction calculus with an additional binary operation between formulas

- \Leftarrow a binary operation \Leftarrow between formulas (called “if”) together with an arrow $(A \Leftarrow B) \wedge B \xrightarrow{\varepsilon_{A,B}} A$ inducing the following *transposition* on arrows:

$$\frac{C \wedge B \xrightarrow{h} A}{C \xrightarrow{\bar{h}} A \Leftarrow B}$$

» Associated proof calculus

One derives two operations on proofs

$$\frac{A \xrightarrow{f} B}{\top \xrightarrow{\overline{f \circ \pi'_{\top, A}}} B \Leftarrow A}$$

$$\frac{\top \xrightarrow{g} B \Leftarrow A}{A \xrightarrow{\varepsilon_{B,A} \circ \langle g \circ A \bullet, 1_A \rangle} B}$$

We denote

$$f_* := \overline{f \circ \pi'_{\top, A}} \quad g^* := \varepsilon_{B,A} \circ \langle g \circ A \bullet, 1_A \rangle$$

f_* is called the **name** of f .

» Deduction theorem

Ring like construction

Given a positive intuitionistic calculus \mathcal{D} , assuming the proof $\top \xrightarrow{x} A$, one gets a new positive intuitionistic calculus $\mathcal{D}(x)$ with the same formulas as \mathcal{D} and where the proofs, called **polynomials**, are freely generated using the induced operators on proofs (inference and derived rules), like $\langle -, - \rangle$, \wedge , $(-)^*$ and $(-)_*$.

Deduction theorem on proofs

With every proof $B \xrightarrow{\varphi(x)} C$ in $\mathcal{D}(x)$ from the *assumption* $\top \xrightarrow{x} A$, there exists an associated proof $A \wedge B \xrightarrow{f} C$ in \mathcal{D} not depending on x .

» Other deduction systems

One can go further and define

- * **intuitionistic** propositional calculus (adding falsehood and disjunction)
- * classical **propositional calculus** (adding the excluded middle)

» Deduction systems as categories

We can fully recover a category structure from a deduction system by adding back the missing axioms as equivalence relation between proofs.

More precisely, the equality between proofs is decided modulo the following identities

- * $f \circ 1_A = f$, for any object A and arrow f with domain A
- * $1_A \circ f = f$, for any object A and arrow f with codomain A
- * $(f \circ g) \circ h = f \circ (g \circ h)$, for any composable arrows f, g, h

» Conjunction calculus as cartesian category

Conjunction calculus can be regarded as a cartesian category by restricting further the equality between proofs modulo the following identities:

- * $f = A \bullet$, for any $A \xrightarrow{f} \top$ (now \top becomes a terminal object)
- * for all $C \xrightarrow{f} A$, $C \xrightarrow{g} B$, $C \xrightarrow{h} A \wedge B$
 - * $\pi_{A,B} \circ \langle f, g \rangle = f$
 - * $\pi'_{A,B} \circ \langle f, g \rangle = g$
 - * $\langle \pi_{A,B} \circ h, \pi'_{A,B} \circ h \rangle = h$
- * This turns the conjunction into a product

» Positive intuitionistic calculus as cartesian closed category

We restrict further the equalities between proofs modulo the following identities:

for all $C \wedge B \xrightarrow{h} A$ and $C \xrightarrow{k} A \Leftarrow B$

$$* \varepsilon_{A,B} \langle \bar{h} \circ \pi_{C,B}, \pi'_{C,B} \rangle = h$$

$$* \overline{\varepsilon_{A,B} \circ \langle k \circ \pi_{C,B}, \pi'_{C,B} \rangle} = k$$

These identities make the “if” binary operation \Leftarrow into an **exponential**, so $B \Leftarrow A$ defines B^A which satisfies all the properties of a (right) adjunction for the product functor.

» Polynomial category

Let \mathcal{A} denote the cartesian closed category obtained from a positive intuitionistic calculus \mathcal{D} .

The **polynomial category** $\mathcal{A}[x]$ is defined as the cartesian closed category obtained from the associated positive intuitionistic calculus $\mathcal{D}(x)$ assuming $\top \xrightarrow{x} A$.

Remark: It can be shown that $\mathcal{A}[x]$ is isomorphic to a **Kleisli category** but we won't be detailing such construction (unfortunately).

» A deduction theorem over categories

For any polynomial $\varphi(x) : B \rightarrow C$ in $\mathcal{A}[x]$ there is a unique arrow $f : A \times B \rightarrow C$ in \mathcal{A} such that $f \circ \langle x \circ B \bullet, 1_B \rangle = \varphi(x)$. (The equality here is between equivalence classes by construction of \mathcal{A} and $\mathcal{A}[x]$.)

This says that polynomials have very **special canonical form**, very much like $a_0 + a_1X + a_2X^2 + \dots$ is the canonical form of polynomials over $k[X]$.

But nothing says that the arrow f is simple!

» Functional completeness

For any polynomial $\varphi(x) : T \rightarrow B$ in $\mathcal{A}[x]$, where $x : T \rightarrow \mathcal{A}$ is an assumption in \mathcal{A} , there is a unique arrow $f : T \rightarrow B^A$ in \mathcal{A} such that $\varepsilon_{B,A} \circ \langle f, x \rangle = \varphi(x)$ (again the equality is over equivalence classes). We denote f by $\lambda_{x:A} \varphi(x)$.

Yes, this will be the λ abstraction in the typed λ -calculus.

λ -calculus

» Untyped λ-calculus

Combinatory logic

The **pure λ-calculus** is a formal language. Its words, called **λ-terms** are defined inductively

$$t ::= x \mid t \wr t' \mid \lambda_x.t$$

where the (total) binary operator \wr , called **application** and the binder λ (over variables), called **λ-abstraction** satisfy the following axioms:

- (β) $(\lambda_x.\varphi(x)) \wr a = \varphi(a)$, whenever no free occurrence in a becomes bound in $\varphi(a)$; we say x is **substituted** by a , or a is **substitutable** for x .
- (η) $\lambda_x.(f \wr x) = f$, whenever f is **independent** from x (i.e. if x occurs in f it must be bound).

A term is **closed** if it contains no free variables.

» α -renaming

Equivalence relation

Terms are considered equal up to renaming their bound variables.

This defines a **congruence** relation (equality over equivalence classes).

For instance $\lambda x.y \equiv \lambda z.y \not\equiv \lambda y.y$

» Typed λ -calculus

A **typed λ -calculus** is a formal language consisting of

- * a class of types
- * a class of terms for each type

The class of types

- * has some **basic types** (like \top , or N for natural numbers)
- * is **closed** under products and exponentials: for any types A and B , $A \times B$ and B^A are also types.

The class of terms is **freely generated**

- * from variables of certain types
- * term forming operations: pairing $\langle -, - \rangle$, projections π, π' , evaluation $\varepsilon_{A,B}$, and λ -abstraction

» Translations

A **translation** is a morphism over λ -calculi $\phi : \mathcal{L} \rightarrow \mathcal{L}'$:

- * $\phi(A)$ is a type of \mathcal{L}' for any type A of \mathcal{L}
- * $\phi(1) = 1$, $\phi(A \times B) = \phi(A) \times \phi(B)$ etc.
- * for every arrow $a : 1 \rightarrow A$ in \mathcal{L} , $\phi(a) : 1 \rightarrow \phi(A)$ in \mathcal{L}'
- * if a is closed, then $\phi(a)$ is closed

This defines a category $\lambda\text{-Calc}$ of λ -calculi.

» Internal language of cartesian closed categories

Let \mathcal{A} denote a cartesian closed categories.

The **internal language** $L(\mathcal{A})$ of \mathcal{A} is defined by

- * types are the formulas of \mathcal{A}
- * terms of type A are polynomial expressions $\varphi(x) : 1 \rightarrow A$ in the polynomial category $\mathcal{A}[x]$ where $x : 1 \rightarrow B$ (typed variables).
Notice that the domain for terms is the terminal object 1. Thus any arrow in the polynomial category is not a term, but its **name** is (cf. slide 14).
- * (We need a natural numbers object to have multiple variables.)

Terms are “**ordinary elements**” of types.

» Curry-Howard-Lambek isomorphism

The internal language construction defines a functor

$$L : \mathbf{Cart} \rightarrow \lambda\text{-Calc}$$

We can also **generate** a cartesian closed category from a λ -calculus. This defines a functor $C : \lambda\text{-Calc} \rightarrow \mathbf{Cart}$

Curry-Howard-Lambek isomorphism

$$\lambda\text{-Calc} \cong \mathbf{Cart}$$

(The equivalence still stands if one add a (weak) natural number object and state the correspondence with such object added in in both the language and the cartesian closed category.)

» This is only the big bang

You have already seen a lot up to this point.

But that's just the beginning!

- * The construction of $\mathcal{A}[x]$ using Kleisli categories
- * Monads and algebraic theories
- * reduction and bounded (strongly normalizing) λ -terms (coherence problem)
- * C-monoidal categories and untyped λ -calculus
- * ...

To be continued ...

» References

- * Joachim Lambek, Philip J. Scott. Introduction to Higher-Order Categorical Logic. 1986
- * Steve Awodey, Category Theory. 2009